
## Lesson 8.5 Exercises, pages 680-688

## Α

**3.** For each graph of y = f(x), draw vertical lines to represent the vertical asymptotes, if they exist, of the graph of  $y = \frac{1}{f(x)}$ .



 $\begin{array}{c|c}
 & y \\
\hline
 & x \\
\hline
 & y = f(x) \\
\hline
 & -6 \\
\hline
 & -8 \\
\hline
\end{array}$ 

The *x*-intercepts of the graph of y = f(x) are 2 and 6.

So, the graph of  $y = \frac{1}{f(x)}$ has vertical asymptotes at x = 2 and x = 6.


- The x-intercept of the graph of y = f(x) is -2. So, the graph of  $y = \frac{1}{f(x)}$  has a vertical asymptote at x = -2.
- **4.** For each graph in question 3, identify the values of x for which the graph of  $y = \frac{1}{f(x)}$  is above the x-axis, and for which it is below the x-axis.
  - a) Above the x-axis: 2 < x < 6</li>Below the x-axis: x < 2 or x > 6
- b) Below the *x*-axis:  $x \in \mathbb{R}, x \neq -2$ Above the *x*-axis: never

**5.** On each graph from question 3, sketch the graph of  $y = \frac{1}{f(x)}$ .

a) y = f(x)  $y = \frac{1}{f(x)}$   $y = \frac{1}{f(x)}$ 

The x-axis is a horizontal asymptote. Plot points where the lines y = 1 and y = -1 intersect the graph. These points are common to both graphs. The graph of y = f(x) has vertex (4, 4), so point  $\left(4, \frac{1}{4}\right)$  lies on  $y = \frac{1}{f(x)}$ .


Since the graph has 2 vertical asymptotes, it has Shape 3.




The x-axis is a horizontal asymptote. Plot points where the line y = -1 intersects the graph. These points are common to both graphs. Since the graph has one vertical asymptote, it has Shape 2.

**6.** Match each graph of y = f(x) to the corresponding graph of  $y = \frac{1}{f(x)}$ . What features of the graphs did you use to make your decisions?

a) -6 -4 -2 0 y = f(x)



i)  $2 \frac{y}{y} = \frac{1}{f(x)}$   $-4 -2 \qquad 2 \qquad 4$ 



The graph that corresponds to graph a is graph ii. The graph of y = f(x) in part a has 2 x-intercepts, so the graph of its reciprocal function has 2 vertical asymptotes.

The graph that corresponds to graph b is graph i. The graph of y = f(x) in part b has no x-intercepts, so the graph of its reciprocal function has no vertical asymptotes.

## В

**7.** How many vertical asymptotes does the graph of each reciprocal function have? Identify the equation of each vertical asymptote.

a) 
$$y = \frac{1}{-(x+3)^2}$$
  
 $-(x+3)^2 = 0$  when  $x = -3$ . So, the graph of  $y = \frac{1}{-(x+3)^2}$  has 1 vertical asymptote,

x = -3.

b) 
$$y = \frac{1}{(x-2)(x+6)}$$
  
 $(x-2)(x+6) = 0$  when  $x = 2$  or  $x = -6$ . So, the graph of  $y = \frac{1}{(x-2)(x+6)}$  has 2 vertical asymptotes,  $x = 2$  and  $x = -6$ .

c) 
$$y = \frac{1}{x^2 + x - 6}$$
  
 $x^2 + x - 6 = (x + 3)(x - 2)$   
 $(x + 3)(x - 2) = 0$  when  
 $x = -3$  or  $x = 2$ . So, the  
graph of  $y = \frac{1}{x^2 + x - 6}$  has  
2 vertical asymptotes,  $x = -3$   
and  $x = 2$ .

$$y = \frac{1}{x^2 + x - 6}$$

$$x^2 + x - 6 = (x + 3)(x - 2)$$

$$(x + 3)(x - 2) = 0 \text{ when}$$

$$x = -3 \text{ or } x = 2. \text{ So, the}$$

$$\text{graph of } y = \frac{1}{x^2 + x - 6} \text{ has}$$


$$2 \text{ vertical asymptotes.}$$

$$d) y = \frac{1}{-3x^2 - 9}$$


$$-3x^2 - 9 = -3(x^2 + 3)$$

$$\text{Since } x^2 + 3 \text{ cannot be 0, the graph of } y = \frac{1}{-3x^2 - 9} \text{ has no vertical asymptotes.}$$

**8.** On the graph of each quadratic function y = f(x), sketch a graph of the reciprocal function  $y = \frac{1}{f(x)}$ . Identify the vertical asymptotes, if they exist. Explain your strategies.




The graph of y = f(x) has no x-intercepts, so the graph of  $y = \frac{1}{f(x)}$  has no vertical asymptotes and has Shape 1. Horizontal asymptote: y = 0Points (0, -4), (-1, -5), and (1, -5) lie on y = f(x), so points (0, -0.25), (-1, -0.2), and (1, -0.2) lie on  $y = \frac{1}{f(x)}$ .



The graph of y = f(x) has 2 x-intercepts, so the graph of  $y = \frac{1}{f(x)}$  has 2 vertical asymptotes, x = -3 and x = -1, and has Shape 3. Horizontal asymptote: y = 0Plot points where the lines y = 1 and y = -1intersect the graph of y = f(x). These points are common to both graphs. The graph of y = f(x)has vertex (-2, -2), so point (-2, -0.5)lies on  $y = \frac{1}{f(x)}$ 

c)



The graph of y = f(x) has 1 *x*-intercept, so the graph of  $y = \frac{1}{f(x)}$  has 1 vertical


$$y = \frac{1}{f(x)}$$
 it is a vertical asymptote,  $x = 3$ , and

asymptote, x = 3, and has Shape 2.

Horizontal asymptote: y = 0Plot points where the line y = 1 intersects the graph of y = f(x). These points are

common to both graphs.

d)



The graph of y = f(x) has no

*x*-intercepts, so the graph of  $y = \frac{1}{f(x)}$ 

has no vertical asymptotes and has Shape 1.


Horizontal asymptote: y = 0

Points (3, 2), (1, 4), and (5, 4) lie on y = f(x), so points (3, 0.5), (1, 0.25),

and (5, 0.25) lie on  $y = \frac{1}{f(x)}$ .


**9.** On the graph of each reciprocal function  $y = \frac{1}{f(x)}$ , sketch a graph of the quadratic function y = f(x).

a)




The graph has one vertical asymptote, x = -5, so the graph of y = f(x) has vertex (-5, 0). The line y = -1 intersects the graph at 2 points that are common to both graphs.

b)




The graph has 2 vertical asymptotes, x = -1 and x = 1, so the graph of y = f(x) has x-intercepts -1 and 1.

The point  $\left(0, -\frac{1}{4}\right)$  is on the line of symmetry, so (0, -4) is the vertex of y = f(x).





The graph has no vertical asymptote, so the graph of y = f(x) has no x-intercept. The point (-6, 1) is on the line of symmetry, so (-6, 1) is the vertex of y = f(x). Points (-7, 0.5) and (-5, 0.5) lie on y = f(x), so points (-7, 2) and (-5, 2) lie on  $y = \frac{1}{f(x)}$ .



The graph has one vertical asymptote, x = 5, so the graph of y = f(x) has vertex (5, 0). The line y = 1 intersects the graph at 2 points that are common to both graphs.

**10.** The graphs of the reciprocal functions  $y = \frac{1}{f(x)}$  and  $y = \frac{1}{g(x)}$  are shown below.






- a) Identify whether each function y = f(x) and y = g(x) is linear or quadratic. How did you decide?
  - i) y = f(x) is quadratic. I recognize the shape of the graph as that of the reciprocal of a quadratic function with 1 x-intercept.
- ii) y = g(x) is linear. I recognize the shape of the graph as that of the reciprocal of a linear function.
- **b**) On the graph of each reciprocal function, sketch a graph of the related linear or quadratic function. What strategy did you use each time?
  - i) The graph has one vertical asymptote, x = 4, so the graph of y = f(x) has vertex (4, 0). The line y = -1 intersects the graph at 2 points that are common to both graphs.
- ii) Vertical asymptote is about x = -1.8, so graph of y = g(x) has x-intercept -1.8. Mark points at y = 1 and y = -1 on graph of  $y = \frac{1}{g(x)}$ , then draw a line through these points for the graph of y = g(x).

**11.** Graph each pair of functions on the same grid. Explain your strategies.

**a)** 
$$y = 4(x + 2)^2$$
 and  $y = \frac{1}{4(x + 2)^2}$ 

The graph of  $y = 4(x + 2)^2$  opens up, has vertex (-2, 0), and x-intercept -2. The graph of  $y = \frac{1}{4(x + 2)^2}$  has vertical asymptote, x = -2 and horizontal asymptote y = 0. Plot points where the line y = 1 intersects the graph of  $y = 4(x + 2)^2$ . These points are common to both graphs. The graph of the reciprocal function has Shape 2.



**b**) 
$$y = -2x^2 - 3$$
 and  $y = \frac{1}{-2x^2 - 3}$ 

The graph of  $y = -2x^2 - 3$  opens down, with vertex (0, -3), so the graph has no x-intercepts. The graph of  $y = \frac{1}{-2x^2 - 3}$ 

has no vertical asymptotes; the horizontal asymptote is y = 0. Points (0, -3), (1, -5), and (-1, -5) lie on  $y = -2x^2 - 3$ .

So, points 
$$\left(0, -\frac{1}{3}\right)$$
, (1,  $-0.2$ ), and

$$(-1, -0.2)$$
 lie on  $y = \frac{1}{-2x^2 - 3}$ . The graph

of the reciprocal function has Shape 1.

$$y = \frac{1}{-2x^2 - 3} \frac{2}{-4} \frac{y}{-2} \frac{x}{4} \frac{y}{-4} \frac{y}{-2} \frac{y}{4} \frac{y}{-2} \frac$$


c) 
$$y = 2x^2 - 4x - 6$$
 and  $y = \frac{1}{2x^2 - 4x - 6}$ 

The graph of  $y = 2x^2 - 4x - 6$ , or y = 2(x - 3)(x + 1) opens up, has x-intercepts 3 and -1, and vertex (1, -8). The graph of

 $y = \frac{1}{2x^2 - 4x - 6}$  has vertical asymptotes x = 3 and x = -1, and a horizontal asymptote y = 0. Plot points where the lines y = 1and y = -1 intersect the graph of y = 2(x - 3)(x + 1). These points are common to both graphs. Point (1, -8) lies on  $y = 2x^2 - 4x - 6$ ,




The graph of the reciprocal function has Shape 3.



**12.** A quadratic function and its reciprocal are graphed on a grid. The horizontal lines y = 1 and y = -1 do not intersect either graph. Sketch a possible graph of the quadratic function and its reciprocal. Explain your strategy.

If the lines y=1 and y=-1 do not intersect the graph of a quadratic function, the vertex of the graph either lies above the line y=1 and opens up, or it lies below the line y=-1 and opens down. In either case, the graph of the quadratic function has no x-intercepts and the graph of the corresponding reciprocal function has no vertical asymptotes. Here is the graph of  $y=x^2+3$  and  $y=\frac{1}{x^2+3}$ .



- **13.** A reciprocal function has the form  $y = \frac{1}{ax^2 + b}$ ,  $a \ne 0$ ,  $b \ne 0$ . How can you determine the number of asymptotes of the graph of  $y = \frac{1}{ax^2 + b}$  when:
  - **a**) both *a* and *b* are positive?

Look at the quadratic function  $y = ax^2 + b$ . When both a and b are positive, the graph opens up and its vertex is above the x-axis. So, the graph of  $y = ax^2 + b$  has no x-intercepts and the graph of  $y = \frac{1}{ax^2 + b}$  has no vertical asymptotes.

**b**) both *a* and *b* are negative?

Look at the quadratic function  $y = ax^2 + b$ . When both a and b are negative, the graph opens down and its vertex is below the x-axis. So, the graph of  $y = ax^2 + b$  has no x-intercepts and the graph of  $y = \frac{1}{ax^2 + b}$  has no vertical asymptotes.

**c)** *a* and *b* have opposite signs?

Look at the quadratic function  $y = ax^2 + b$ . When a and b have opposite signs, the graph either opens up with its vertex below the x-axis, or it opens down with its vertex above the x-axis. So, the graph of  $y = ax^2 + b$  has 2 x-intercepts and the graph of  $y = \frac{1}{ax^2 + b}$  has 2 vertical asymptotes.

## C

**14.** The reciprocal function  $y = \frac{1}{px^2 + (2p + 1)x + p}$  has one vertical asymptote. Show that  $p = -\frac{1}{4}$ .

If the reciprocal function has one vertical asymptote, then the related quadratic function has one x-intercept; that is, the quadratic equation  $px^2 + (2p + 1)x + p = 0$  has equal roots.

The equation has equal roots when  $b^2 - 4ac = 0$ .

Substitute: 
$$b = 2p + 1$$
,  $a = p$ ,  $c = p$ 

$$(2p + 1)^2 - 4(p)(p) = 0$$

$$4p^2 + 4p + 1 - 4p^2 = 0$$

$$4p + 1 = 0$$

$$p=-\frac{1}{4}$$

**15.** Determine the values of *k* for which the reciprocal function

$$y = \frac{1}{x^2 + kx + 4}$$
 has:

a) no vertical asymptotes

The reciprocal function has no vertical asymptotes, so the related quadratic function has no x-intercepts; that is,  $x^2 + kx + 4 = 0$  has no real roots. The equation has no real roots when  $b^2 - 4ac < 0$ .

Substitute: 
$$a = 1$$
,  $b = k$ ,  $c = 4$ 

$$k^2 - 4(1)(4) < 0$$

$$k^2 - 16 < 0$$

$$k^2 < 16$$

$$-4 < k < 4$$

**b**) one vertical asymptote

The reciprocal function has one vertical asymptote, so the related quadratic function has one x-intercept; that is,  $x^2 + kx + 4 = 0$  has equal roots. The equation has equal roots when  $b^2 - 4ac = 0$ .

Substitute: 
$$a = 1$$
,  $b = k$ ,  $c = 4$ 

$$k^2 - 4(1)(4) = 0$$

$$k^2-16=0$$

$$k^2 = 16$$

$$k = 4 \text{ or } k = -4$$

c) two vertical asymptotes

The reciprocal function has two vertical asymptotes, so the related quadratic function has two x-intercepts; that is,  $x^2 + kx + 4 = 0$ has 2 real roots. The equation has 2 real roots when  $b^2 - 4ac > 0$ . Substitute: a = 1, b = k, c = 4

$$k^2 - 4(1)(4) > 0$$

$$k^2 - 16 > 0$$

$$k^2 > 16$$

$$k^{-} > 16$$
  
  $k < -4 \text{ or } k > 4$ 

8.5 Graphing Reciprocals of Quadratic Functions—Solutions

- **16.** Determine the equation of each quadratic function y = f(x) you graphed in question 9. Describe your strategies.
  - a) The graph has vertex (-5, 0) and passes through the point (-6, -1). So, the equation has the form:  $y = a(x + 5)^2$ Substitute: x = -6, y = -1 $-1 = a(-6 + 5)^2$ -1 = a

The equation of the quadratic function is  $y = -(x + 5)^2$ .

b) The graph has vertex (0, -4) and passes through the point (1, 0). So, the equation has the form:  $y = ax^2 - 4$ Substitute: x = 1, y = 0

$$0 = a(1)^2 - 4$$

4 = a

The equation of the quadratic function is  $y = 4x^2 - 4$ .

c) The graph has vertex (-6, 1) and passes through the point (-5, 2). So, the equation has the form:  $y = a(x + 6)^2 + 1$ 

```
Substitute: x = -5, y = 2
2 = a(-5 + 6)^2 + 1
```

1 = a

The equation of the quadratic function is  $y = (x + 6)^2 + 1$ .

d) The graph has vertex (5, 0) and passes through the point (4, 1).

So, the equation has the form:  $y = a(x - 5)^2$ 

Substitute: x = 4, y = 1

 $1 = a(4 - 5)^2$ 

1 = a

The equation of the quadratic function is  $y = (x - 5)^2$ .