

Lesson 4.2 Exercises, pages 278–284

A

3. Given $f(x) = x^3$ and $g(x) = x^2 + 1$, write an explicit equation for each combination.

a) $h(x) = f(x) + g(x)$

$$h(x) = x^3 + x^2 + 1$$

b) $d(x) = f(x) - g(x)$

$$d(x) = x^3 - x^2 - 1$$

c) $p(x) = f(x) \cdot g(x)$

$$p(x) = x^3(x^2 + 1)$$

d) $q(x) = \frac{f(x)}{g(x)}$

$$q(x) = \frac{x^3}{x^2 + 1}$$

4. For each function $h(x)$ below, write explicit equations for $f(x)$ and $g(x)$ so that:

i) $h(x)$ is the sum $f(x) + g(x)$

ii) $h(x)$ is the difference $f(x) - g(x)$

a) $h(x) = x^2 + 3x - 4$

b) $h(x) = x^3 - x^2 + 8$

Sample answers:

i) $h(x) = x^2 + (3x - 4)$

$$f(x) = x^2 \text{ and } g(x) = 3x - 4$$

i) $h(x) = (x^3 - x^2) + 8$

$$f(x) = x^3 - x^2 \text{ and } g(x) = 8$$

ii) $h(x) = x^2 - (-3x + 4)$

$$f(x) = x^2 \text{ and } g(x) = -3x + 4$$

ii) $h(x) = x^3 - (x^2 - 8)$

$$f(x) = x^3 \text{ and } g(x) = x^2 - 8$$

B

5. Use $f(x) = 2x - 4$ and $g(x) = -x + 2$.

a) Write an explicit equation for $h(x)$.

i) $h(x) = f(x) + g(x)$

$$h(x) = 2x - 4 + (-x + 2)$$

$$h(x) = x - 2$$

ii) $h(x) = g(x) + f(x)$

$$h(x) = -x + 2 + 2x - 4$$

$$h(x) = x - 2$$

iii) $h(x) = f(x) - g(x)$

$$h(x) = 2x - 4 - (-x + 2)$$

$$h(x) = 3x - 6$$

iv) $h(x) = g(x) - f(x)$

$$h(x) = -x + 2 - (2x - 4)$$

$$h(x) = -3x + 6$$

v) $h(x) = f(x) \cdot g(x)$

$$h(x) = (2x - 4)(-x + 2)$$

$$h(x) = -2x^2 + 8x - 8$$

vi) $h(x) = g(x) \cdot f(x)$

$$h(x) = (-x + 2)(2x - 4)$$

$$h(x) = -2x^2 + 8x - 8$$

- b) For part a, compare the answers to parts i and ii; parts iii and iv; and parts v and vi. Explain the results.

The answers to parts i and ii are the same because addition is commutative. The answers to parts iii and iv are opposites because subtraction is not commutative. The answers to parts v and vi are the same because multiplication is commutative.

6. Given that $f(x) = x^2 - 4$, $g(x) = 2x - 1$, and $h(x) = 3 - x^3$, write an explicit equation for $k(x)$, then state its domain.

a) $k(x) = f(x) + g(x) + h(x)$ b) $k(x) = f(x) - g(x) + h(x)$

$$k(x) = x^2 - 4 + 2x - 1 + 3 - x^3$$

$$k(x) = -x^3 + x^2 + 2x - 2$$

This is a cubic function; its domain is: $x \in \mathbb{R}$

$$k(x) = x^2 - 4 - (2x - 1) + 3 - x^3$$

$$k(x) = -x^3 + x^2 - 2x$$

This is a cubic function; its domain is: $x \in \mathbb{R}$

c) $k(x) = f(x) + g(x) \cdot h(x)$ d) $k(x) = g(x) \cdot f(x) - h(x)$

$$k(x) = x^2 - 4 + (2x - 1)(3 - x^3)$$

$$k(x) = x^2 - 4 + 6x - 2x^4 - 3 + x^3$$

$$k(x) = -2x^4 + x^3 + x^2 + 6x - 7$$

This is a quartic function; its domain is: $x \in \mathbb{R}$

$$k(x) = (2x - 1)(x^2 - 4) - (3 - x^3)$$

$$k(x) = 2x^3 - 8x - x^2 + 4 - 3 + x^3$$

$$k(x) = 3x^3 - x^2 - 8x + 1$$

This is a cubic function; its domain is: $x \in \mathbb{R}$

7. Use the function $k(x) = x^2 - 3x - 28$.

- a) Write explicit equations for three functions $f(x)$, $g(x)$, and $h(x)$ so that $k(x) = f(x) + g(x) + h(x)$.

Sample response:

$$k(x) = x^2 - 3x - 28$$

$$k(x) = (x^2) + (-3x) + (-28)$$

$$f(x) = x^2; g(x) = -3x; h(x) = -28$$

- b) Write explicit equations for two functions $f(x)$ and $g(x)$ so that $k(x) = f(x) \cdot g(x)$.

Sample response:

$$k(x) = x^2 - 3x - 28$$

$$\text{Factor: } k(x) = (x - 7)(x + 4)$$

$$f(x) = x - 7; g(x) = x + 4$$

8. For each function $h(x)$ below, write explicit equations for $f(x)$ and $g(x)$ so that:

- i) $h(x)$ is the sum $f(x) + g(x)$
- ii) $h(x)$ is the difference $f(x) - g(x)$
- iii) $h(x)$ is the product $f(x) \cdot g(x)$
- iv) $h(x)$ is the quotient $\frac{f(x)}{g(x)}$

a) $h(x) = x^2$

b) $h(x) = \sqrt{x}$

Sample response:

i) Subtract and add the same term.

$$h(x) = x^2 - x + x$$

$$f(x) = x^2 - x; g(x) = x$$

ii) Add and subtract the same term.

$$h(x) = x^2 + x - x$$

$$f(x) = x^2 + x; g(x) = x$$

iii) Write x^2 as a product.

$$h(x) = (x)(x)$$

$$f(x) = x; g(x) = x$$

iv) Multiply and divide x^2 by the same non-zero expression.

$$h(x) = \frac{x^2(x^2 + 1)}{x^2 + 1}$$

$$f(x) = x^2(x^2 + 1);$$

$$g(x) = x^2 + 1$$

i) Subtract and add the same term.

$$h(x) = \sqrt{x} - x + x$$

$$f(x) = \sqrt{x} - x; g(x) = x$$

ii) Add and subtract the same term.

$$h(x) = \sqrt{x} + x - x$$

$$f(x) = \sqrt{x} + x; g(x) = x$$

iii) Multiply \sqrt{x} by a term that is equal to 1.

$$h(x) = \sqrt{x} \left(\frac{2}{2} \right)$$

$$f(x) = 2\sqrt{x}; g(x) = \frac{1}{2}$$

iv) Multiply and divide \sqrt{x} by the same non-zero expression.

$$h(x) = \sqrt{x} \left(\frac{2 + x^2}{2 + x^2} \right)$$

$$f(x) = (2 + x^2)\sqrt{x};$$

$$g(x) = 2 + x^2$$

9. Use $f(x) = |x - 4|$ and $g(x) = x^2$.

a) State the domain and range of $f(x)$ and of $g(x)$.

$f(x)$ is an absolute value function; the domain is $x \in \mathbb{R}$ and the range is $y \geq 0$. $g(x)$ is a quadratic function whose graph has vertex $(0, 0)$ and opens up; the domain is $x \in \mathbb{R}$, and the range is $y \geq 0$.

b) Given $h(x) = f(x) + g(x)$, write an explicit equation for $h(x)$, then determine its domain and range.

$$h(x) = |x - 4| + x^2$$

Since the domains of $f(x)$ and $g(x)$ are equal, then the domain of $h(x)$ is $x \in \mathbb{R}$. Use technology to graph the function; the minimum value is 3.75 at $x = 0.5$, so the range is $y \geq 3.75$.

- c) Given $d(x) = f(x) - g(x)$, write an explicit equation for $d(x)$, then determine its domain and range.

$$d(x) = |x - 4| - x^2$$

Since the domains of $f(x)$ and $g(x)$ are equal, then the domain of $d(x)$ is $x \in \mathbb{R}$. Use technology to graph the function; the maximum value is 4.25 at $x = -0.5$, so the range is $y \leq 4.25$.

10. Use $f(x) = x^3 - x$ and $g(x) = \frac{1}{x + 3}$.

- a) State the domain and range of $f(x)$ and of $g(x)$.

$f(x)$ is a cubic function; the domain is $x \in \mathbb{R}$ and the range is $y \in \mathbb{R}$.

$g(x)$ is a reciprocal function; the domain is $x \neq -3$, and the range is $y \neq 0$.

- b) Given $h(x) = f(x) + g(x)$, write an explicit equation for $h(x)$, then determine its domain and range.

$$h(x) = x^3 - x + \frac{1}{x + 3}$$

The domain of $h(x)$ is the set of values of x that are common to the domains of $f(x)$ and $g(x)$, so the domain is $x \neq -3$. Use technology to graph the function; the approximate range is $y \leq -34.5$ or $y \geq -14.2$.

- c) Given $p(x) = f(x) \cdot g(x)$, write an explicit equation for $p(x)$, then determine its domain and range.

$$p(x) = \frac{x^3 - x}{x + 3}$$

The domain of $p(x)$ is the set of values of x that are common to the domains of $f(x)$ and $g(x)$, so the domain is $x \neq -3$. Use technology to graph the function; the range is $y \in \mathbb{R}$.

11. Use $f(x) = \sqrt{x + 2}$ and $g(x) = |x - 2|$.

- a) State the domain and range of $f(x)$ and of $g(x)$.

$f(x)$ is a square root function; the domain is $x \geq -2$ and the range is $y \geq 0$. $g(x)$ is an absolute value function; the domain is $x \in \mathbb{R}$, and the range is $y \geq 0$.

- b) Given $p(x) = f(x) \cdot g(x)$, write an explicit equation for $p(x)$, then determine its domain and range.

$$p(x) = \sqrt{x + 2} \cdot |x - 2|$$

The domain of $p(x)$ is all values of x that are common to the domains of $f(x)$ and $g(x)$, so the domain is $x \geq -2$. Use technology to graph the function; the range is $y \geq 0$.

- c) Given $q(x) = \frac{f(x)}{g(x)}$, write an explicit equation for $q(x)$, then determine its domain and range.

$$q(x) = \frac{\sqrt{x+2}}{|x-2|}$$

The domain of $q(x)$ is restricted to those values of x for which $|x-2| \neq 0$ and for which $\sqrt{x+2}$ is defined, so the domain is $x \geq -2$, $x \neq 2$. Use technology to graph the function; the range is $y \geq 0$.

12. a) When asked to write $f(x) = x^2$ as the quotient of two functions, a student wrote $f(x) = \frac{x^3}{x}$. Is this correct? Justify your answer.

No, the answer is incorrect because the domain of the new function has the restriction $x \neq 0$, which the original function did not have.

- b) If your answer to part a is no, write $f(x) = x^2$ as a quotient of two functions.

Multiply and divide the function by a non-zero expression, such as $(x^2 + 1)$.

A possible function is: $f(x) = \frac{x^2(x^2 + 1)}{x^2 + 1}$

13. Consider the functions: $f(x) = (x + 3)^2$ and $g(x) = \frac{x - 2}{x + 3}$

Given $p(x) = f(x) \cdot g(x)$, write an explicit equation for $p(x)$, then determine its domain and range.

$$p(x) = (x + 3)^2 \left(\frac{x - 2}{x + 3} \right), \text{ or } p(x) = (x + 3)(x - 2), x \neq -3$$

$$p(x) = x^2 + x - 6, x \neq -3$$

The domain is $x \neq -3$. Use technology to graph the function; the range is $y \geq -6.25$.

14. Consider the function $g(x) = 4$ and any function $f(x)$. Predict how the graph of each function below will be a transformation image of $y = f(x)$. Use graphing technology to check.

a) $y = f(x) + g(x)$

b) $y = f(x) - g(x)$

The function $g(x)$ is a horizontal line with y -intercept 4.

When $g(x)$ is added to $f(x)$, the graph of $y = f(x)$ will be translated 4 units up.

When $g(x)$ is subtracted from $f(x)$, the graph of $y = f(x)$ will be translated 4 units down.

c) $y = f(x) \cdot g(x)$

d) $y = \frac{f(x)}{g(x)}$

When $f(x)$ is multiplied by $g(x)$, the graph of $y = f(x)$ will be stretched vertically by a factor of 4.

When $f(x)$ is divided by $g(x)$, the graph of $y = f(x)$ will be compressed vertically by a factor of $\frac{1}{4}$.

15. When each function $h(x)$ below is evaluated at $x = a$, its value is 0.

What do you know about the values of $f(a)$ and $g(a)$?

a) $h(x) = f(x) + g(x)$

Substitute: $x = a, h(a) = 0$

$$0 = f(a) + g(a)$$

$$f(a) = -g(a)$$

b) $h(x) = f(x) - g(x)$

Substitute: $x = a, h(a) = 0$

$$0 = f(a) - g(a)$$

$$f(a) = g(a)$$

c) $h(x) = f(x) \cdot g(x)$

Substitute: $x = a, h(a) = 0$

$$0 = f(a) \cdot g(a)$$

$$f(a) = 0, \text{ or } g(a) = 0, \text{ or both}$$

d) $h(x) = \frac{f(x)}{g(x)}$

Substitute: $x = a, h(a) = 0$

$$0 = \frac{f(a)}{g(a)}$$

$$f(a) = 0 \text{ and } g(a) \neq 0$$

16. Given $f(x) = \sqrt{x}$ and $g(x) = \sqrt{2-x}$, determine an explicit equation for each function, then state its domain.

a) $h(x) = f(x) + g(x)$

$$h(x) = \sqrt{x} + \sqrt{2-x}$$

For $f(x)$, $x \geq 0$ and for $g(x)$,

$x \leq 2$, so the domain of $h(x)$ is:

$$0 \leq x \leq 2$$

b) $d(x) = f(x) - g(x)$

$$d(x) = \sqrt{x} - \sqrt{2-x}$$

For $f(x)$, $x \geq 0$ and for $g(x)$,

$x \leq 2$, so the domain of $d(x)$ is:

$$0 \leq x \leq 2$$

c) $p(x) = f(x) \cdot g(x)$

$$p(x) = \sqrt{x} \cdot \sqrt{2-x}$$

For $f(x)$, $x \geq 0$ and for $g(x)$,

$x \leq 2$, so the domain of $p(x)$ is:

$$0 \leq x \leq 2$$

d) $q(x) = \frac{f(x)}{g(x)}$

$$q(x) = \frac{\sqrt{x}}{\sqrt{2-x}}$$

For $f(x)$, $x \geq 0$ and for $g(x)$,

$x \leq 2$, but since $g(x)$ is in the

denominator, $x \neq 2$

The domain of $q(x)$ is: $0 \leq x < 2$

C

17. Consider the function: $f(x) = \frac{x^2 - 3x + 4}{x - 1}$

a) Determine the domain and the approximate range of $f(x)$.

Since the denominator cannot be 0, the domain is: $x \neq 1$

Use technology to graph the function.

It has a minimum point at approximately (2.4, 1.8) and a maximum

point at approximately (-0.4, -3.8).

So, the range is approximately $y \leq -3.8$ or $y \geq 1.8$.

b) Determine explicit equations for $g(x)$, $h(x)$, and $k(x)$ so that

$$f(x) = g(x) + \frac{h(x)}{k(x)}.$$

Sample response: Use synthetic division to determine: $(x^2 - 3x + 4) \div (x - 1)$

$$\begin{array}{r|rrr} 1 & 1 & -3 & 4 \\ & & 1 & -2 \\ \hline & 1 & -2 & 2 \end{array}$$

The function can be written as:

$$f(x) = x - 2 + \frac{2}{x - 1}$$

So, $g(x) = x - 2$; $h(x) = 2$; and $k(x) = x - 1$

18. Is it possible to combine $f(x) = \sqrt{x}$ with a second function $g(x)$ to get a new function whose domain is all real numbers? Justify your answer.

No, when two functions are combined, the domain of the new function is the set of values of x that are common to the two functions that were combined. Since the domain of \sqrt{x} is $x \geq 0$, then the domain of the new function cannot be all real numbers.