

REVIEW, pages 560–566

6.1

1. Determine the value of each trigonometric ratio. Use exact values where possible; otherwise write the value to the nearest thousandth.

a) $\tan(-45^\circ)$
 $= -1$

b) $\cos 600^\circ$
 $= \cos 240^\circ$
 $= -\frac{1}{2}$

c) $\sec(-210^\circ)$
 $= \frac{1}{\cos(-210^\circ)}$
 $= \frac{1}{\cos 150^\circ}$
 $= -\frac{2}{\sqrt{3}}$

d) $\sin 765^\circ$
 $= \sin 45^\circ$
 $= \frac{1}{\sqrt{2}}$

e) $\cot 21^\circ$
 $= \frac{1}{\tan 21^\circ}$
 $\doteq 2.605$

f) $\csc 318^\circ$
 $= \frac{1}{\sin 318^\circ}$
 $\doteq -1.494$

2. To the nearest degree, determine all possible values of θ for which $\cos \theta = 0.76$, when $-360^\circ \leq \theta \leq 360^\circ$.

Since $\cos \theta$ is positive, the terminal arm of angle θ lies in Quadrant 1 or 4.

The reference angle is: $\cos^{-1}(0.76) \doteq 41^\circ$

For the domain $0^\circ \leq \theta \leq 360^\circ$:

In Quadrant 1, $\theta \doteq 41^\circ$

In Quadrant 4, $\theta \doteq 360^\circ - 41^\circ$, or approximately 319°

For the domain $-360^\circ \leq \theta \leq 0^\circ$:

In Quadrant 1, $\theta \doteq -360^\circ + 41^\circ$, or approximately -319°

In Quadrant 4, $\theta \doteq -41^\circ$

6.2

3. As a fraction of π , determine the length of the arc that subtends a central angle of 225° in a circle with radius 3 units.

Arc length: $\frac{225}{360}(2\pi)(3) = \frac{15}{4}\pi$

6.3

4. a) Convert each angle to degrees. Give the answer to the nearest degree where necessary.

i) $\frac{5\pi}{3}$

$$= \frac{5(180^\circ)}{3}$$
$$= 300^\circ$$

ii) $-\frac{10\pi}{7}$

$$= -\frac{10(180^\circ)}{7}$$
$$\doteq -257^\circ$$

iii) 4

$$= 4\left(\frac{180^\circ}{\pi}\right)$$
$$\doteq 229^\circ$$

b) Convert each angle to radians.

i) 150°

$$= 150 \left(\frac{\pi}{180} \right)$$
$$= \frac{5\pi}{6}$$

ii) -240°

$$= -240 \left(\frac{\pi}{180} \right)$$
$$= -\frac{4\pi}{3}$$

iii) 485°

$$= 485 \left(\frac{\pi}{180} \right)$$
$$= \frac{97\pi}{36}$$

5. In a circle with radius 5 cm, an arc of length 6 cm subtends a central angle. What is the measure of this angle in radians, and to the nearest degree?

Angle measure is: $\frac{\text{arc length}}{\text{radius}} = \frac{6}{5}$

$$= 1.2$$

In degrees, $1.2 = 1.2 \left(\frac{180^\circ}{\pi} \right)$

$$\doteq 69^\circ$$

The angle measure is 1.2 radians or approximately 69° .

6. A race car is travelling around a circular track at an average speed of 120 km/h. The track has a diameter of 1 km. Visualize a line segment joining the race car to the centre of the track. Through what angle, in radians, will the segment have rotated in 10 s?

In 1 s, the car travels: $\frac{120}{60 \cdot 60} \text{ km} = \frac{1}{30} \text{ km}$

So, in 10 s, the car travels: $\frac{10}{30} \text{ km} = \frac{1}{3} \text{ km}$

Angle measure is: $\frac{\text{arc length}}{\text{radius}} = \frac{\frac{1}{3}}{1}$

$$= \frac{1}{3}$$

In 10 s, the segment will have rotated through an angle of $\frac{1}{3}$ radian.

7. Determine the value of each trigonometric ratio. Use exact values where possible; otherwise write the value to the nearest thousandth.

a) $\sin \frac{\pi}{3}$

$$= \frac{\sqrt{3}}{2}$$

b) $\cos \frac{5\pi}{6}$

$$= -\frac{\sqrt{3}}{2}$$

c) $\sec \left(-\frac{\pi}{2} \right)$

$$= \frac{1}{\cos \left(-\frac{\pi}{2} \right)}$$

which is undefined

d) $\tan \frac{15\pi}{4}$

$$= \tan \frac{3\pi}{4}$$
$$= -1$$

e) $\csc 5$

$$= \frac{1}{\sin 5}$$
$$\doteq -1.043$$

f) $\cot (-22.8)$

$$= \frac{1}{\tan (-22.8)}$$
$$\doteq -0.954$$

8. P(3, -1) is a terminal point of angle θ in standard position.

a) Determine the exact values of all the trigonometric ratios for θ .

Let the distance between the origin and P be r .

Use: $x^2 + y^2 = r^2$ Substitute: $x = 3, y = -1$

$$9 + 1 = r^2$$

$$r = \sqrt{10}$$

$$\sin \theta = -\frac{1}{\sqrt{10}} \quad \csc \theta = -\sqrt{10} \quad \cos \theta = \frac{3}{\sqrt{10}}$$

$$\sec \theta = \frac{\sqrt{10}}{3} \quad \tan \theta = -\frac{1}{3} \quad \cot \theta = -3$$

b) To the nearest tenth of a radian, determine possible values of θ in the domain $-2\pi \leq \theta \leq 2\pi$.

The terminal arm of angle θ lies in Quadrant 4.

The reference angle is: $\tan^{-1}\left(\frac{1}{3}\right) = 0.3217\dots$

So, $\theta = -0.3217\dots$

The angle between 0 and 2π that is coterminal with $-0.3217\dots$ is:

$$2\pi - 0.3217\dots = 5.9614\dots$$

Possible values of θ are approximately: 6.0 and -0.3

6.4

9. Use graphing technology to graph each function below for $-2\pi \leq x \leq 2\pi$, then list these characteristics of the graph: amplitude, period, zeros, domain, range, and the equations of the asymptotes.

a) $y = \sin x$

The amplitude is 1. The period is 2π . The zeros are $0, \pm\pi, \pm 2\pi$. The domain is $-2\pi \leq x \leq 2\pi$. The range is $-1 \leq y \leq 1$. There are no asymptotes.

b) $y = \cos x$

The amplitude is 1. The period is 2π . The zeros are $\pm\frac{\pi}{2}, \pm\frac{3\pi}{2}$.

The domain is $-2\pi \leq x \leq 2\pi$. The range is $-1 \leq y \leq 1$. There are no asymptotes.

c) $y = \tan x$


There is no amplitude. The period is π . The zeros are $0, \pm\pi, \pm 2\pi$.

The domain is $x \neq \pm\frac{\pi}{2}, \pm\frac{3\pi}{2}$. The range is $y \in \mathbb{R}$. The equations of the asymptotes are $x = \pm\frac{\pi}{2}$ and $x = \pm\frac{3\pi}{2}$.

6.5

10. On the same grid, sketch graphs of the functions in each pair for $0 \leq x \leq 2\pi$, then describe your strategy.

a) $y = \sin x$ and $y = \sin\left(x - \frac{\pi}{4}\right)$




For the graph of $y = \sin x$, I used the completed table of values from Lesson 6.4.

The horizontal scale is 1 square to $\frac{\pi}{4}$ units, because the phase shift is $\frac{\pi}{4}$.

I then shifted several points $\frac{\pi}{4}$ units right and joined the points to get the graph of $y = \sin\left(x - \frac{\pi}{4}\right)$.

b) $y = \cos x$ and $y = \frac{3}{2} \cos x$



For the graph of $y = \cos x$, I used the completed table of values from Lesson 6.4.


I multiplied every y -coordinate by 1.5, plotted the new points, then

joined them to get the graph of $y = \frac{3}{2} \cos x$.

6.6

11. a) Graph $y = \frac{1}{2} \sin 3\left(x + \frac{\pi}{6}\right) + 2$ for $-2\pi \leq x \leq 2\pi$.

Explain your strategy.




Sample response: I graphed $y = \frac{1}{2} \sin 3x$, shifted several points $\frac{\pi}{6}$ units left and 2 units up, then joined the points to get the graph of $y = \frac{1}{2} \sin 3\left(x + \frac{\pi}{6}\right) + 2$.

- b) List the characteristics of the graph you drew.

The amplitude is $\frac{1}{2}$. The period is $\frac{2\pi}{3}$. There are no zeros.

The domain is $-2\pi \leq x \leq 2\pi$. The range is $\frac{3}{2} \leq y \leq \frac{5}{2}$.

12. An equation of the function graphed below has the form $y = a \cos b(x - c) + d$. Identify the values of a , b , c , and d in the equation, then write an equation for the function.



Sample response: The equation of the centre line is $y = \frac{1}{2}$, so the vertical translation is $\frac{1}{2}$ unit up and $d = \frac{1}{2}$.

The amplitude is: $\frac{3 - (-2)}{2} = \frac{5}{2}$, so $a = \frac{5}{2}$

Choose the x -coordinates of two adjacent maximum points,

$-\frac{\pi}{3}$ and $\frac{11\pi}{3}$. The period is: $\frac{11\pi}{3} - \left(-\frac{\pi}{3}\right) = 4\pi$

So, b is: $\frac{2\pi}{4\pi} = \frac{1}{2}$


To the left of the y -axis, the cosine function begins its cycle at $x = -\frac{\pi}{3}$, so a possible phase shift is $-\frac{\pi}{3}$, and $c = -\frac{\pi}{3}$.

Substitute for a , b , c , and d in: $y = a \cos b(x - c) + d$

An equation is: $y = \frac{5}{2} \cos \frac{1}{2}\left(x + \frac{\pi}{3}\right) + \frac{1}{2}$

6.7

13. A water wheel has diameter 10 m and completes 4 revolutions each minute. The axle of the wheel is 8 m above a river.



- a) The wheel is at rest at time $t = 0$ s, with point P at the lowest point on the wheel. Determine a function that models the height of P above the river, h metres, at any time t seconds. Explain how the characteristics of the graph relate to the given information.

The time for 1 revolution is 15 s.

At $t = 0, h = 3$

At $t = 7.5, h = 13$

The graph begins at $(0, 3)$, which is a minimum point.

The first maximum point is at $(7.5, 13)$.

The next minimum point is after 1 cycle and it has coordinates $(15, 3)$.

The position of the first maximum is known, so use a cosine function:

$$h(t) = a \cos b(t - c) + d$$


The constant in the equation of the centre line of the graph is the height of the axle above the river, so its equation is: $h = 8$; and this is also the vertical translation, so $d = 8$

The amplitude is one-half the diameter of the wheel, so $a = 5$

The period is the time for 1 revolution, so $b = \frac{2\pi}{15}$

A possible phase shift is: $c = 7.5$

An equation is: $h(t) = 5 \cos \frac{2\pi}{15}(t - 7.5) + 8$



- b) Use technology to graph the function. Use this graph to determine:

- i) the height of P after 35 s

Graph: $Y = 5 \cos \frac{2\pi}{15}(X - 7.5) + 8$

Determine the Y-value when $X = 35$.

After 35 s, P is 10.5 m high.

ii) the times, to the nearest tenth of a second, in the first 15 s of motion that P is 11 m above the river

Graph: $Y = 5 \cos \frac{2\pi}{15}(X - 7.5) + 8$ and $Y = 11$

Determine the Y-coordinates of the first two points of intersection.

P is 11 m above the river after approximately 5.3 s and 9.7 s.